

# OKLAHOMA | 2017 Healthy Housing Fact Sheet

Unhealthy conditions found in hazardous housing can lead to lead poisoning, asthma, respiratory illness, cancer, and unintentional injuries, resulting in missed school days and poor school performance for children, as well as missed work days for parents. To protect the health of Oklahoma's families and prevent continued increases in associated healthcare costs and societal consequences, full federal funding is needed for critical programs and services:

	OK received funding since 2015?
<b>CDC's Healthy Homes and Lead Poisoning Prevention Program</b>	YES
<b>CDC's National Asthma Control Program</b>	NO
<b>CDC's Environmental Health Tracking Network</b>	NO
<b>HUD's Office of Lead Hazard Control and Healthy Homes</b>	NO
<b>HUD's Community Development Block Grants (CDBG)</b>	YES
<b>HUD's HOME Investment Partnerships Program</b>	YES
<b>EPA's Lead Categorical Grants</b>	YES
<b>EPA's Drinking Water Revolving Fund</b>	YES
<b>HHS' Maternal and Child Health Block Grants</b>	YES
<b>HHS' Low Income Home Energy Assistance Program (LIHEAP)</b>	YES
<b>DOE's Weatherization Assistance Program</b>	YES

Full funding to federal programs such as these will help to address many of the risks and burdens facing the families and residents of Oklahoma, including:



In Oklahoma, **26% of children live in households with a high housing cost burden**, and 22% of children live in poverty.



Approximately **10% of adults and 12% of children have current asthma** in Oklahoma (2014).



**55% of Oklahoma housing was built before 1978** and is likely to contain lead-based paint.



In 2012, **total charges for asthma-related hospitalizations in Oklahomans were \$76.4 million**, \$18.5 million of which is attributable to **1,928 asthma-related hospitalizations for Oklahoman children under 15 years of age**.



In 2015, **1,213 Oklahoma children tested had an elevated blood lead level (5 µg/dL or more); 164 of them had blood lead levels of 10 µg/dL or more**.



On average, **11 individuals die annually from carbon monoxide exposure in Oklahoma** (2011-2015).



**Every week, 9 older adults die and 124 are hospitalized** from fall-related injuries in Oklahoma.



**9 counties** in Oklahoma are categorized as having **"moderate potential" for indoor radon levels above the EPA action level**.